Spider BAS

COMPOSITION: Spirotetramat 15% (w/v)
Complementary materials up to 100% (w/v)

INDICATIONS:

CROP	PEST	Application Rate ml/20 liter	Interval between applications (minimum)	Number of applications/ Season	Pre-Harvest Interval (PHI)
Brassicas: Broccoli, Cabbage, Cauliflower, Turnip	Aphids, Whiteflies	7.5-12	7 days	*2	3 days
Forest Trees	Aphids, Scale insects	12-24	14 days	2	3 days
Citrus: Clementine, Grapefruit, Kumquat, Lemon, Mandarin, Tangerine, Orange, Pomelo	Aphids, Thrips, Citrus mite, Scale insects, Mealybugs, Leaf miners, Whiteflies	18-24	21 days	2	3 days
Fruiting Vegetables: Eggplant, Physalis, Okra, Hot Pepper, Sweet Pepper, Tomato	Aphids, Broad mite, Psyllid, Tomato rust mite, Whiteflies	7.5-12	7 days	2	3 days
Leafy Vegetables: Arugula, Artichoke, Celery, Chicory, Lettuce, Purslane, Parsley, Chard, Spinach	Aphids, Whiteflies	7.5-12	7 days	2	3 days
Legumes: Peas, Lupine, Beans, Broad Beans, Chickpeas, Lentils	Aphids, Whiteflies	7.5-12	7 days	2	Beans Fresh: 1-3 days Dry: 7 days
Pome Fruits: Apple, Pear, Loquat, Quince	Aphids, Rust mite, Mealybugs, Pear psylla, Whiteflies	12-21	14 days	**3	7 days
Roots and Tubers: Ginger, Potato, Sweet Potato, Turmeric	Aphids, Psyllid, Whiteflies	9-12	7 days	2	7 days
Vine Fruits: Grapes, Kiwi	Grape galls, Mealybugs, Grape phylloxera Whiteflies,	12-18	30 days	2	7 days
Stone Fruits: Apricot, Cherry, Nectarine, Plum, Peach	Aphids, Mealybugs, Scale insects, Whiteflies	12-21	14 days	1	7 days
Sugar Beet	Bean aphid, Whiteflies	11-21.6	14 days	2	28 days
Nuts: Almond, Cashew, Pistachio, Walnut	Aphids, Mealybugs, Phylloxera, Scale insects, Whiteflies	12-21	14 days	3	7 days
Tropical Fruits: Mango, Guava, Avocado	Aphids, Avocado thrips, Mealybugs, Scale insects, Whiteflies	20-24	14 days	2	3 days

Properties: An insecticide that is primarily effective against immature stages of the target pests. It also inhibits reproduction in adult target pests. The insecticide is translocated through the phloem and xylem to all plant tissues, including newly developing shoots, leaves, and roots.

Re-entry interval (for treated fields): Entry into treated areas is prohibited for 24 hours after application.

Product image



Toxic agricultural pesticide

Group	23	INSECTICIDE

Antidote

No specific antidote is available; treat the affected individual symptomatically according to the clinical signs and symptoms presented.

STORAGE

Store the pesticide in its original, tightly sealed container in a cool, dry, secure, and well-ventilated area, away from direct sunlight.

Keep the pesticide away from animals, food, feed, seeds, and fertilizers.

Packaging

100/250/500/1000 ml

ISO CERTIFIED : ISO 45001__ ISO 14001__ISO 9001__ISO 22000__HACCP

صنع في الأردن
Made in Jordan

سبيدر بس

المكونات: سبيروتيترامات ١٥٪ (و/ح)
مواد مكملة حتى ١٠٠٪ (و/ح)

طريقة الإستعمال:

المحصول	الآفة	معدل التطبيق مل / 20 لتر	الحد الأدنى للفاصل الزمني بين الرشة والأخرى/ يوم	عدد الرشات المسموحة بالموسم	فترة الأمان (PHI) / يوم
البروكلي، ملفوف، الزهرة، اللفت	المن، الذبابة البيضاء	12-7.5	7	*2	3
الأشجار الحرجية	المن، الحشرات القشرية	24-12	14	2	3
الحمضيات: (كلمنتينا، الجريب فروت، الكمكوات، الليمون، مندلينا، اليوسفي، البرتقال، بوملي)	المن، التربس، حلم الحمضيات، الحشرات القشرية، البق الدقيقي، صانعات الأنفاق، الذبابة البيضاء	24-18	21	2	3
الخضروات المثمرة: الباذنجان، الحرنكش، الباميه، الفلفل الحار، الفلفل الحلو، البذورة	المن، الحلم العريض، البسيلا حلم صدأ البذورة، الذبابة البيضاء	12-7.5	7	2	3
الخضروات الورقية: الجرجير، الخرشوف الشوكي، الكرفس، الهندباء، الخس، البقلة، البقدونس، السلق، السبانخ	المن، الذبابة البيضاء	12-7.5	7	2	3
البقوليات: البازلاء، التربس، الفاصولياء، الفول العريض، الحمص، العدس	المن، الذبابة البيضاء	12-7.5	7	2	الفاصولياء الطازجة: 1-3 الجافة: 7
التفاح، الاجاص، اسكدنيا ، السفرجل	المن، حلم الصدأ، البق الدقيقي، بسيلا الإجاص، الذبابة البيضاء	21-12	14	**3	7
الزنجبيل، البطاطا، البطاطا الحلوة، الكركم	المن، البسيلا، الذبابة البيضاء	12 9	7	2	7
العنب، الكيوي	تورم العنب، البق الدقيقي الفلوكسيرا، الذبابة البيضاء	18-12	30	2	7
المشمش، الكرز ،نكتارين، الخوخ، الدراق	المن، البق الدقيقي، الحشرات القشرية، الذبابة البيضاء	21-12	14	1	7
الشمندر السكري	من الفاصولياء، الذبابة البيضاء	21.6-11	14	2	28
اللوز ،الكاجو ،الفستق الحلبي، الجوز	المن، البق الدقيقي، الفلوكسيرا، الحشرات القشرية، الذبابة البيضاء	21-12	14	3	7
الفواكه الاستوائية: المنجا، الجوافة، الأفوكادو	المن، تربس الأفوكادو، البق الدقيقي، الحشرات القشرية، الذبابة البيضاء	24-20	14	2	3

الخواص : مبيد حشري فعال بشكل أساسي ضد الآفات المستهدفة غير الناضجة، كما يعمل على تثبيط تكاثر الآفات المستهدفة البالغة ويتحرك المبيد عبر اللحاء والخشب إلى جميع أنسجة النبات بما في ذلك أنسجة البراعم والأوراق والجذور الجديدة.

فترة تحريم دخول الحقل المرشوش: ثباتية محلول الرش: لا يسمح بدخول المناطق المعالجة خلال ٢٤ ساعة.

صورة المنتج



مبيد زراعي سام

INSECTICIDE	23	Group

الجرعة المضادة

لا يوجد جرعة مضادة، يعالج المصاب حسب الأعراض الظاهرة.

التخزين

يحفظ المبيد في عبوته الأصلية محكمة الإغلاق في منطقة جافة وباردة وآمنة وجيدة التهوية، بعيداعن اشعة الشمس المباشرة. يحفظ بعيدًا عن الحيوانات والطعام والأعلاف والذور والأسمدة.

التعبئة

١٠٠٠/٥٠٠/٢٥٠/١٠٠ مل

ISO CERTIFIED : ISO 45001__ ISO 14001__ISO 9001__ISO 22000__HACCP

صنع في الأردن
Made in Jordan