Exofeda

Oral solution For treatment and control of lice and mites

COMPOSITION: Each 1 ml contains:
Fluralaner 10 mg

PROPERTIES AND INDICATIONS:
Exofeda is effective acaricide against mites infesting poultry, sheep, goats and cattle. It is a potent inhibitor of parts of the arthropods nervous system. It begins to work within four hours of the mites feeding on treated animal and subsequently stop eggs production by female mites for 15 days after the first dose, disrupting the mite's life cycle.
Exofeda used for the treatment of poultry red mites (Dermanyssus gallina) in pullets, chickens for reproduction and layer hens.

TARGET SPECIES: Chickens (pullets, chickens for reproduction and layer hens), sheep, Goats and Cattle.

DOSAGE AND ADMINISTRATION:
Poultry: For use in drinking water: 0.05 ml of Exofeda per 1 kg of body weight (equivalent to 0.5 mg fluralaner) administered twice, 7 days apart. The complete course of treatment must be administered for a full therapeutic effect.
For each day of treatment, medicated water must be freshly prepared.
If another course of treatment is indicated, the interval between two courses of treatment should be at least 3 months.
Sheep, Goats and Cattle: 1.5 ml per 10 kg of body weight, administered orally using drenching tools.
Do not repeat treatment before 54 days.
Do not mix treated and untreated animals.

Product image



WITHDRAWAL PERIOD

Meat: 14 days.
Eggs: None.
Milk: Not for use in animals producing milk for human consumption.

STORAGE

Store in a dry place away from direct heat (below than 30°C).

Packaging

100/1000 ml

ISO CERTIFIED : ISO 45001__ ISO 14001__ ISO 9001__ ISO 22000__ HACCP

صنع في الأردن
Made in Jordan

(للإستعمال البيطري فقط)
(For Veterinary Use Only)

إكسوفدا

محلول فموي للعلاج والسيطرة على القمل والعث

المكونات: كل ١ مل يحتوي :
فلورالانير ١٠ ملغم

الخصائص والإستعمالات:
اكسوفدا هو مبيد فعال ضد العث الذي يصيب الدواجن والأغنام والماعز والأبقار، وهو مثبط قوي لأجزاء من الجهاز العصبي للمفصليات، يبدأ مفعوله خلال أربع ساعات من بدء تغذية العث على الحيوان المُعالج وبالتالي يوقف إنتاج البيض من إناث السوس لمدة ١٥ يومًا بعد أول جرعة مما يؤدي الى تعطيل دورة حياة العث.
يُستخدم اكسوفدا لعلاج عث الدواجن الأحمر (ديرمانيسوس جالينا) في الدجاجات الصغيرة، ودجاج التكاثر، والدجاج البياض.

الحيوانات المستهدفة: الدجاج (الدجاج الصغير، دجاج التكاثر والدجاج البياض) والاغنام والماعز والابقار

الجرعة وطريقة الإستخدام :
الدواجن: للاستخدام في مياه الشرب: يُعطى ٠,٠٥ مل من اكسوفدا لكل ١ كغ من وزن الجسم (ما يعادل ٠,٥ ملغم فلورالانير) مرتين، بفاصل ٧ أيام.
يجب تناول جرعة العلاج كاملةً للحصول على تأثير علاجي كامل.
في كل يوم من أيام العلاج، يجب تحضير الماء المعالج حديثًا.
في حال الحاجة إلى جرعة علاجية أخرى، يجب ألا تقل الفترة الفاصلة بين جرعتين عن ٣ أشهر.
الأغنام ،الماعز والابقار: ١,٥ مل لكل ١٠ كغم من وزن الجسم تعطى عن طريق الفم باستخدام أدوات التجريع.
لا تكرر العلاج قبل ٥٤ يومًا.
لا تخلط الحيوانات المعالجة مع غير المعالجة.

صورة المنتج



فترة السحب
اللحوم: ١٤ يوم.
البيض: لا يوجد
الحليب: لا يستخدم للحيوانات المنتجة للحليب للاستهلاك البشري

التخزين
يحفظ في مكان جاف بعيداً عن الحرارة المباشرة
(اقل من ٣٠ درجة مئوية).

التعبئة
١٠٠ / ١٠٠٠ مل

(للإستعمال البيطري فقط)
(For Veterinary Use Only)

صنع في الأردن
Made in Jordan

ISO CERTIFIED : ISO 45001__ ISO 14001__ISO 9001__ISO 22000__HACCP